

# JENNIEFER KHO, M.D.

Surgery of Hand, Orthopaedic Surgery

## Main Office

4849 Van Nuys Blvd Suite #217  
Sherman Oaks, CA 91403

T: 818-902-2800

F: 818-782-8979

10640 Riverside Dr  
North Hollywood, CA 91602

T: 818-955-9898

23043 Lyons Ave.  
Santa Clarita, CA 91321

T:661-254-8163

## POST-OPERATIVE INSTRUCTIONS

1. Keep the hand ELEVATED as much as possible until swelling is minimal.
2. MOVE the fingers, unless your fingers are in a splint.
3. Keep the dressing DRY. Put a plastic bag or glove, and tape, over the dressing during showers until sutures are removed (usually 2 weeks).
4. If you have had a carpal tunnel release or trigger finger release, you may remove the surgical dressing in 5 days, and place a BAND-AID over the incision. Otherwise, keep the dressing ON unless instructed otherwise,
5. You may ICE the hand/fingers as needed to reduce swelling and pain. Ice for no more than 20 MINUTES at a time.
6. If you were prescribed a narcotic, use it sparingly for moderate to severe pain. Otherwise, an NSAID and Tylenol are sufficient for pain relief.
7. Numbness, tingling, and swelling is very common after hand surgery. If there is severe swelling in the hand, ELEVATE the hand first. If not better after a few hours, then call the office.
8. Nausea/vomiting is very common after anesthesia. Stay HYDRATED and avoid narcotics if you have this symptom. Constipation is also a common side effect of narcotics.
9. Call the office for a follow-up appointment as directed by Dr. Kho (1-2 weeks).